

HEUCHERA INDIAN SUMMER



FLAMINGO

Holland Inc.



Bilberry



Blackberry Ruffle



Boysenberry



Coralberry




Cranberry

- // Outstanding heat and sun tolerance
- // Great for combos
- // Compact foliage
- // Zone 4-9
- // Can be grown with minimal shade
- // Gorgeous color range



Gajiberry




Gooseberry



Huckleberry




Mulberry




Orange Berry Ruffle



Raspberry



Sloeberry



Tayberry

Transplanting

Remove liner carefully from trays so as not to damage roots. Plant liner slightly above soil line being mindful to not bury crown when transplanting. Keep soil surface of liner level with surface of finished pot.

Container Sizes and Crop Times

- Quart/4" from 72-cell liner: 1ppp. 8-10 weeks after transplant.
- Gallon/6" from 72-cell liner: 1ppp. 12-14 weeks after transplant.
- 8" from 72-cell liner: 3ppp. 12-14 weeks after transplant.

Growing Media

Plant into a well-drained media with a pH: 5.8-6.2.

Temperature

Night temperature 50-55°F (10-13°C).

Day temperature 60-65°F (16-18°C).

Vernalization is not required.

Light

Low to moderate light levels of 1,500-2,500 foot candles.

Heuchera is day length neutral.

Water and Feed

Heuchera should be grown using a well balanced fertilizer at a rate of 100-125 ppm.

The media should be allowed to dry slightly between watering and never saturated.

Growth Regulators

In most conditions, PGR application will not be necessary.

Insects and Diseases

Pythium and Rhizoctonia can occur if plants are overwatered. Heuchera can generally be grown free of insects. Aphids, mealybugs, slugs, spider mites, thrips and whiteflies may appear occasionally. Preventive measures are typically not necessary.

